



FLWEMS Paramedics Neonatal & Pediatric Protocol for the Management of:  
**ANAPHYLAXIS & ALLERGIC REACTIONS**

**Indications**

To outline the paramedic care and management of the neonatal/pediatric patient with signs & symptoms of acute altered mental status changes.

**Procedures**

1. Secure an airway as outlined in FLWEMS Paramedics Neonatal & Pediatric Protocol for the Management of Airway & Ventilation and administer supplemental **Oxygen** as needed. Intubate neurologically depressed patient to prevent aspiration.
2. Apply cardiac monitor, pulse oximetry and/or capnometry as needed.
3. Utilize the "Broslow Tape" system for procedure and medication administration guidelines.
4. Establish IV or IO access.
5. IV infusion **0.9% NaCL** titrated to SBP appropriate for age group.
6. Consider **Epinephrine** .01mg/kg SQ (maximum of 0.3mg). May repeat every 15 minutes as needed. May administer prior to the establishment of IV/IO access if patient is in severe distress.
7. Consider **Diphenhydramine** (Benadryl) 1-2mg/kg IVP.
8. Consider **Albuterol** (Proventil) 2.5mg via NEB for wheezing.
9. Consider **Methylprednisolone** (Solu-Medrol) 1-2mg/kg IVP.
10. Transport to appropriate Emergency Department.
11. Contact Medical Control for further orders as needed.

**CAIRA/Chemical Surety Considerations**

None

**Triage Considerations**

Refer to S.T.A.R.T. Triage Protocol

**END OF SOP – NOTHING FOLLOWS**